

Vaccinations	Before Pregnancy	During Pregnancy	After Pregnancy	Type of Vaccine
<b>Tdap</b>	<b>Yes,</b> If Recommended	<b>Yes,</b> must get vaccinated between 27 and 36 weeks of gestation	<b>Yes,</b> If not given previously	Inactivated
<b>Tetanus/Diphtheria Td</b>	<b>Yes,</b> If Recommended	<b>Yes,</b> If Recommended	<b>Yes,</b> If not given previously	Toxoid
<b>Influenza IIV</b>	<b>Yes</b>	<b>Yes</b>	<b>Yes</b>	Inactivated
<b>Hepatitis A</b>	<b>Yes,</b> If Recommended	<b>Yes,</b> If Recommended	<b>Yes,</b> If Recommended	Inactivated
<b>Hepatitis B</b>	<b>Yes,</b> If Recommended	<b>Yes,</b> If Recommended	<b>Yes,</b> If Recommended	Inactivated
<b>Human Papillomavirus (HPV)</b>	<b>Yes,</b> if Recommended, through 26yrs of age	<b>NO</b>	<b>Yes,</b> if Recommended, through 26yrs of age	Inactivated
<b>MMR</b>	<b>Yes,</b> if recommended, avoid conception for 4 weeks	<b>NO</b>	<b>Yes,</b> if Recommended	Live
<b>Meningococcal</b>	If recommended	If recommended	If recommended	Inactivated
<b>Pneumococcal</b>	If recommended	If recommended	If recommended	Inactivated
<b>Varicella</b>	<b>Yes,</b> if recommended, avoid conception for 4 weeks	<b>NO</b>	<b>Yes,</b> if recommended, give immediately postpartum if susceptible	Live
<b>COVID -19</b>	<b>Yes,</b> If recommended	<b>Yes,</b> If recommended	<b>Yes,</b> If recommended	Inactivated

If you are pregnant or planning for pregnancy, Then you must talk with your Doctor before getting vaccinated to avoid future complications. Doctors in your Country, Region can recommend right vaccinations on basis of your specific health conditions.